





Three new species of *Fomitiporia* (Hymenochaetales, Basidiomycota) from tropical dry forests in Ecuador and Peru

Jian Chen^{1*}, Josef Vlasák^{2*}, Kai-Yue Luo¹, Yuan Yuan¹

¹ State Key Laboratory of Efficient Production of Forest Resources, School of Ecology and Nature Conservation, Beijing Forestry University, Beijing 100083, China

² Biology Centre of the Academy of Sciences of the Czech Republic, České Budějovice, Czech Republic

Corresponding authors: Kai-Yue Luo (kaiyue0415@bjfu.edu.cn); Yuan Yuan (yuanyuan1018@bjfu.edu.cn)



This article is part of:

**Exploring the Hidden Fungal Diversity:
Biodiversity, Taxonomy, and Phylogeny of
Saprobic Fungi**

Edited by Samantha C. Karunarathna,
Danushka Sandaruwan Tennakoon, Ajay
Kumar Gautam

Academic editor:

Samantha C. Karunarathna

Received: 28 March 2025

Accepted: 14 May 2025

Published: 10 June 2025

Citation: Chen J, Vlasák J, Luo K-Y, Yuan Y (2025) Three new species of *Fomitiporia* (Hymenochaetales, Basidiomycota) from tropical dry forests in Ecuador and Peru. MycoKeys 118: 267–287. <https://doi.org/10.3897/mycokeys.118.154175>

Copyright: © Jian Chen et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

Three new species in Hymenochaetaceae are illustrated and described from tropical Ecuador and Peru based on morphology and phylogenetic analyses. The species *Fomitiporia robustiformis* is characterized by perennial, pileate basidiocarps; ungulate pilei; brown and glabrous pileal surface; broad and rounded margin; yellowish-brown pore surface; circular to angular pores measuring 6–7 per mm; thick and entire dissepiments; a dimittic hyphal system; the presence of fusoid cystidioles; the presence of hymenial setae; and globose basidiospores measuring $5\text{--}6.4 \times 4.7\text{--}6.1 \mu\text{m}$. *Fomitiporia roseo-bubalina* is characterized by annual to perennial, resupinate to pileate basidiocarps; ungulate to triquetrous pilei; grayish and glabrous pileal surface; blunt margin; pale gray and glossy pore surface; circular to angular pores measuring 4–5 per mm; thick and entire dissepiments; a dimittic hyphal system; the presence of hymenial setae and cystidioles; and globose basidiospores measuring $5.3\text{--}6.7 \times 5\text{--}6.2 \mu\text{m}$. *Fomitiporia triqueter* is characterized by perennial, pileate, sessile basidiocarps; triquetrous pilei with sharp margins; grayish dark and cracked pileal surfaces with concentrically sulcate, pale yellowish brown pore surfaces; circular pores measuring 9–10 per mm; the absence of hymenial setae and cystidioles; the presence of large rhomboid crystals in the hymenium; and globose basidiospores measuring $3.4\text{--}5.5 \times 3.2\text{--}5.4 \mu\text{m}$. The differences between the new species and morphologically similar and phylogenetically related species are also discussed.

Key words: Hymenochaetaceae, phylogeny, taxonomy, wood-decaying fungi

Introduction

Hymenochaetales is a major group of wood-inhabiting fungi, and more than 1,500 species have been reported in the order (Du et al. 2020; Wu et al. 2022a; Wang et al. 2023; Zhou et al. 2023; Dong et al. 2024; Zhao et al. 2024a, 2024b; Liu et al. 2025; Viner 2025; Yang et al. 2025). Hymenochaetaceae is the main family of the order, in which Wu et al. (2022a) made extensive and systematic studies on the family. *Fomitiporia* Murrill (Murrill 1907), as a genus of Hymenochaetaceae typified by *F. langloisii* Murrill, is a worldwide and important genus (Wu et al. 2022b). The genus is rich in species diversity, and many new species were recently described from subtropical and tropical areas (Dai et al. 2021).

* These authors contributed equally to this work.

The largest fungus was found in the genus (Dai and Cui 2011). Some species of the genus are forest pathogens, such as *F. capensis* M. Fisch., M. Cloete, L. Mostert & F. Halleen, *F. hippophaëicola* (H. Jahn) Fiasson & Niemelä, *F. pseudopunctata* (A. David et al.) Fiasson (= *F. mediterranea* M. Fisch.), and *F. torreyae* Y.C. Dai & B.K. Cui (Fischer 2002; Dai et al. 2007; Cloete et al. 2014; Yuan et al. 2023); and some species have medicinal values, such as *F. ellipsoidea* B.K. Cui & Y.C. Dai, *F. hartigii* (Allesch. & Schnabl) Fiasson & Niemelä (\equiv *Phellinus hartigii* (Allesch. & Schnabl) Pat.), *F. punctata* (P. Karst.) Murrill, and *F. robusta* (P. Karst.) Fiasson & Niemelä (\equiv *Phellinus robustus* (P. Karst.) Bourdot & Galzin) (Dai et al. 2009; Wu et al. 2019; Dai 2022, 2023; Zhang et al. 2023, 2024; Ghobad-Nejhad et al. 2024; Shen et al. 2024). In traditional *Phellinus* s.l. (Fischer 1996), *Fomitiporia* has been called the ‘*Phellinus robustus* complex,’ and it was confirmed as a homogeneous genus in Hymenochaetaceae based on molecular data (Wagner and Fischer 2001). The genus is characterized by mostly perennial, pileate or resupinate basidiocarps, homogeneous context, a dimitic hyphal system, hymenial setae rarely present, basidiospores broadly ellipsoid, subglobose or globose, hyaline, thick-walled, smooth, mostly dextrinoid and cyanophilous; on angiosperms and gymnosperms, and causing a white rot (Chen and Cui 2017; Chen et al. 2021; Wu et al. 2022b).

Numerous studies have dealt with *Fomitiporia* in the last 20 years, and species in *Fomitiporia* have been documented to have a higher than presumed phylogenetic and taxonomic diversity (Vlasák and Vlasák 2016; Liu et al. 2018). Because of the similarity in morphology, many new species of *Fomitiporia* have been distinguished by their phylogenetic and geographical characteristics (Chen and Cui 2017). Recently, based on molecular and morphological analyses, more new species were described from the tropics, such as *F. tasmanica* G.M. Gates, X.H. Ji & Jia J. Chen, *F. eucalypti* Y.C. Dai & X.H. Ji, and *F. rondonii* Alves-Silva & Drechsler-Santos (Chen et al. 2021; Wu et al. 2022b; Manawasinghe et al. 2024). About 70 species of *Fomitiporia* are currently reported worldwide (Wu et al. 2022b).

During a study of wood-decaying fungi in South America, three undescribed species matching the concept of *Fomitiporia* were discovered. To understand their taxonomic placement, phylogenetic analysis was carried out based on the internal transcribed spacers (ITS) and the nuclear large subunit ribosomal DNA (nLSU) sequences according to Wu et al. (2022b). Their morphological characteristics were also described and illustrated.

Materials and methods

Morphological studies

The studied specimens were collected from tropical dry forests in Ecuador and Peru (see Results for details of collection and field photographs). All vouchers are deposited in the Fungarium of the Institute of Microbiology, Beijing Forestry University (BJFC), Beijing, China. Morphological characterization (color, texture, taste, and odor of basidiocarps) was conducted based on field notes and voucher specimens. Microscopic analysis follows Chen et al. (2015), and the special color terms follow Petersen (1996). Mushroom sections were examined for microstructures using a Nikon Eclipse 80i compound microscope equipped with

phase contrast optics, with magnification parameters ranging from 100× to 1000×. For measurements and drawings, slide preparations were stained with cotton blue, Melzer's reagent, and 5% potassium hydroxide. Basidiospores were measured from sections cut from the tubes, and 5% of measurements were excluded from each end of the range and are given in parentheses. Abbreviations employed in descriptions: KOH, 5% aqueous potassium hydroxide solution; IKI, Melzer's reagent (IKI[+]: dextrinoid; IKI-: both inamyloid and non-dextrinoid); CB, Cotton Blue affinity (CB+: cyanophilous; CB-: weakly cyanophilous); L, arithmetic average of spore length; W, arithmetic average of spore width; Q, L/W ratios; n, number of basidiospores/measured from given number of specimens.

DNA extraction, amplification, and sequencing

The CTAB-based rapid plant genome extraction kit (Aidlab Biotechnologies Co., Ltd., Beijing) was used to obtain genomic DNA from dried specimens. The primer pair ITS4 and ITS5 was used for amplification of the ITS region (White et al. 1990), while the primer pair LR0R and LR7 (<http://www.biology.duke.edu/fungi/mycolab/primers.htm>) was used to amplify the nLSU region. The PCR procedure for ITS was as follows: initial denaturation at 95 °C for 3 min, followed by 35 cycles at 94 °C for 40 s, 54 °C for 45 s, and 72 °C for 1 min, and a final extension of 72 °C for 10 min. The PCR procedure for nLSU was as follows: initial denaturation at 94 °C for 1 min, followed by 35 cycles at 94 °C for 30 s, 50 °C for 1 min, and 72 °C for 1.5 min, and a final extension of 72 °C for 10 min. The PCR products were purified and sequenced at the Beijing Genomics Institute in China, using the same primers. The newly generated sequences were deposited in the GenBank database.

Phylogenetic analyses

Reference ITS and nLSU sequences from various *Fomitiporia* species were downloaded from GenBank according to Wu et al. (2022b). Sequences of *Neophellinus uncisetus* (Robledo, Urcelay & Rajchenb.) Y.C. Dai, F. Wu, L.W. Zhou, Vlasák & B.K. Cui, obtained from GenBank, were used as the outgroup. All sequences analyzed in this study are listed in Table 1. ITS and nLSU sequences were aligned with BioEdit (Hall 1999) and ClustalX (Thompson et al. 1997). Then two datasets were separately aligned using MAFFT 7.110 (Katoh and Standley 2013) under the G-INS-i option. Ambiguous regions at the start and the end were deleted. Then, alignments were spliced and transformed into formats in Mesquite 3.2 (Maddison and Maddison 2017) and deposited in TreeBASE (<http://www.treebase.org>; accession number 32101; reviewer access URL: <http://purl.org/phylo/treebase/phyloids/study/TB2:S32101?x-access-code=f1ab253962f03e6e486b3b5ce8a04991&format=html>). The phylogenetic analysis was performed using maximum likelihood (ML) and Bayesian inference (BI) methods based on ITS and nLSU datasets (Wu et al. 2022b). The best evolutionary model was estimated using jModelTest (Guindon and Gascuel 2003; Posada 2008) under the Akaike information criterion. ML and BI analyses were conducted using RAXML-HPC2 through the CIPRES Science Gateway, with all parameters set to their default settings (www.phylo.org; Miller et al. 2010). The BI analysis was conducted in two independent runs, each with four

Table 1. Information for the sequences used in this study.

Species name	Sample no.	Location	GenBank accession no.		References
			ITS	nLSU	
<i>Fomitiporia aethiopica</i>	MUCL 44777	Ethiopia	GU478341	AY618204	Amalfi et al. (2010)
<i>Fomitiporia aethiopica</i>	MUCL 44806	Ethiopia	GU461944	AY618202	Amalfi et al. (2010)
<i>Fomitiporia alpina</i>	Dai 15735	China	KX639627	KX639645	Wu et al. (2022b)
<i>Fomitiporia apiahyna</i>	MUCL 51485	Ecuador	GU461962	GU461996	Amalfi et al. (2010)
<i>Fomitiporia atlantica</i>	FLOR 47591	Brazil	KU557529	—	Li et al. (2016)
<i>Fomitiporia atlantica</i>	FLOR 58554	Brazil	KU557528	KU557526	Li et al. (2016)
<i>Fomitiporia australiensis</i>	MUCL 49406	Australia	AY624997	GU462001	Chen et al. (2021)
<i>Fomitiporia baccharidis</i>	MUCL 47756	Argentina	JQ087886	JQ087913	Wu et al. (2022b)
<i>Fomitiporia baccharidis</i>	MUCL 47757	Argentina	JQ087887	JQ087914	Wu et al. (2022b)
<i>Fomitiporia bakeri</i>	MUCL 51098	USA	JQ087874	JQ087901	Wu et al. (2022b)
<i>Fomitiporia bambusarum</i>	ICN 200563	Brazil	MN918544	MN918537	Alves-Silva et al. (2020a)
<i>Fomitiporia bambusarum</i>	ICN 200564	Brazil	MN918545	MN918538	Alves-Silva et al. (2020a)
<i>Fomitiporia bambusipileata</i>	ICN 200559	Brazil	MN918546	MN918539	Alves-Silva et al. (2020a)
<i>Fomitiporia bambusipileata</i>	ICN 200557	Brazil	MN918548	MN918541	Alves-Silva et al. (2020a)
<i>Fomitiporia bannaensis</i>	MUCL 46926	Thailand	KF444682	KF444705	Wu et al. (2022b)
<i>Fomitiporia bannaensis</i>	MUCL 46930	China	KF444683	KF444706	Wu et al. (2022b)
<i>Fomitiporia calkinsii</i>	MUCL 51099	USA	KF444686	KF444709	Wu et al. (2022b)
<i>Fomitiporia calkinsii</i>	MUCL 51398	USA	KF444687	KF444710	Wu et al. (2022b)
<i>Fomitiporia capensis</i>	MUCL 53009	South Africa	JQ087890	JQ087917	Wu et al. (2022b)
<i>Fomitiporia capensis</i>	CMW 48613	South Africa	MH599112	MH599122	Wu et al. (2022b)
<i>Fomitiporia carpineae</i>	Dai 18023	China	MH930812	MH930810	Wu et al. (2022b)
<i>Fomitiporia castilloi</i>	MUCL FG10282	French Guiana	JQ087889	JQ087916	Wu et al. (2022b)
<i>Fomitiporia castilloi</i>	MUCL 53980	French Guiana	JX093786	JX093830	Wu et al. (2022b)
<i>Fomitiporia chilensis</i>	BAFC 52942	Chile	NR_164591	NG_068862	Rajchenberg et al. (2019)
<i>Fomitiporia chilensis</i>	CIEFAPcc586	Chile	MK131090	MK193751	Rajchenberg et al. (2019)
<i>Fomitiporia chilensis</i>	CIEFAPcc587	Chile	MK131094	MK193752	Rajchenberg et al. (2019)
<i>Fomitiporia conyana</i>	FLOR 58547	Brazil	KU663298	KU663270	Alves-Silva et al. (2020b)
<i>Fomitiporia conyana</i>	FLOR 58548	Brazil	KU663299	KU663271	Alves-Silva et al. (2020b)
<i>Fomitiporia cupressicola</i>	MUCL 52489	Mexico	JQ087879	JQ087906	Amalfi et al. (2012)
<i>Fomitiporia cupressicola</i>	MUCL 52490	Mexico	JQ087880	JQ087907	Amalfi et al. (2012)
<i>Fomitiporia deserticola</i>	JV 1209/40-J	USA	KT381634	—	Vlasák and Vlasák (2016)
<i>Fomitiporia deserticola</i>	JV 1209/41-J	USA	KT381633	—	Vlasák and Vlasák (2016)
<i>Fomitiporia deserticola</i>	JV 1209/46	USA	KT381632	—	Vlasák and Vlasák (2016)
<i>Fomitiporia dryophila</i>	MUCL 46380	USA	EF429238	EF429219	Wu et al. (2022b)
<i>Fomitiporia dryophila</i>	MUCL 51144	USA	KF444689	KF444712	Wu et al. (2022b)
<i>Fomitiporia elegans</i>	FURB 44484	Brazil	KU663320	—	Alves-Silva et al. (2020b)
<i>Fomitiporia elegans</i>	FLOR 58556	Brazil	KU663319	KU663293	Alves-Silva et al. (2020b)
<i>Fomitiporia erecta</i>	MUCL 49871	France	GU461939	GU461976	Wu et al. (2022b)
<i>Fomitiporia eucalypti</i>	Dai 18586A	Australia	MH971172	MH971217	Wu et al. (2022b)
<i>Fomitiporia eucalypti</i>	Dai 18682	Australia	MH971171	MH971216	Wu et al. (2022b)
<i>Fomitiporia expansa</i>	MUCL 55026	French Guiana	KJ401031	KJ401032	Amalfi and Decock (2014)
<i>Fomitiporia fissurata</i>	JV 1305/4-J	USA	KT381629	—	Vlasák and Vlasák (2016)
<i>Fomitiporia fissurata</i>	JV 1305/5-J	USA	KT381630	—	Vlasák and Vlasák (2016)
<i>Fomitiporia fissurata</i>	JV 1307/5-J	USA	KT381631	—	Vlasák and Vlasák (2016)
<i>Fomitiporia gabonensis</i>	MUCL 47576	Gabon	GU461971	GU461990	Wu et al. (2022b)
<i>Fomitiporia gaoligongensis</i>	Cui 8261	China	KX639624	KX639642	Chen and Cui (2017)
<i>Fomitiporia gatesiae</i>	Dai 18680	Australia	MH971169	MH971214	Wu et al. (2022b)
<i>Fomitiporia gatesiae</i>	Dai 18681	Australia	MH971170	MH971215	Wu et al. (2022b)
<i>Fomitiporia hartigii</i>	MUCL 53551	Estonia	JX093789	JX093833	Wu et al. (2022b)
<i>Fomitiporia hesleri</i>	01-712/2	-	AY340031	—	Fischer and Binder (2004)
<i>Fomitiporia hesleri</i>	01-77/4	-	AY340026	—	Fischer and Binder (2004)
<i>Fomitiporia hippophaëicola</i>	MUCL 31746	Belgium	GU461945	AY618207	Amalfi et al. (2010)
<i>Fomitiporia hippophaëicola</i>	MUCL 31747	Belgium	GU461946	GU461977	Amalfi et al. (2010)

Species name	Sample no.	Location	GenBank accession no.		References
			ITS	nLSU	
<i>Fomitiporia ignea</i>	TX15	USA	MN108101	MN113927	Brown et al. (2020)
<i>Fomitiporia ignea</i>	TX16	USA	MN108102	MN113928	Brown et al. (2020)
<i>Fomitiporia ivindoensis</i>	MUCL 51311	Gabon	GU461952	GU461979	Amalfi et al. (2010)
<i>Fomitiporia ivindoensis</i>	MUCL 51312	Gabon	GU461951	GU461978	Amalfi et al. (2010)
<i>Fomitiporia lagerstroemiae</i>	Dai 18335	Vietnam	MH930813	MH930811	Wu et al. (2022b)
<i>Fomitiporia langloisii</i>	MUCL 46375	USA	EF429242	EF429225	Wu et al. (2022b)
<i>Fomitiporia maxonii</i>	MUCL 46017	Cuba	EF433559	EF429230	Wu et al. (2022b)
<i>Fomitiporia maxonii</i>	MUCL 46037	Cuba	EF433560	EF429231	Wu et al. (2022b)
<i>Fomitiporia nubicola</i>	FLOR 57850	Brazil	KU663303	NG_073602	Alves-Silva et al. (2020b)
<i>Fomitiporia nubicola</i>	FLOR 57857	Brazil	KU663309	KU663282	Alves-Silva et al. (2020b)
<i>Fomitiporia neotropica</i>	MUCL 53114	French Guiana	JX093792	JX093836	Wu et al. (2022b)
<i>Fomitiporia nobilissima</i>	MUCL 47580	Gabon	GU461966	GU461985	Wu et al. (2022b)
<i>Fomitiporia nobilissima</i>	MUCL 51289	Gabon	GU461965	GU461984	Wu et al. (2022b)
<i>Fomitiporia norbulingka</i>	Cui 9722	China	KU364419	KU364429	Chen et al. (2017)
<i>Fomitiporia norbulingka</i>	Cui 9770	China	KU364420	KU364430	Chen et al. (2017)
<i>Fomitiporia norbulingka</i>	Cui 9766	China	KU364417	KU364427	Chen et al. (2017)
<i>Fomitiporia ovoidospora</i>	Dai 18283	Vietnam	MH971167	MH971212	Wu et al. (2022b)
<i>Fomitiporia ovoidospora</i>	Dai 18349	Vietnam	MH971168	MH971213	Wu et al. (2022b)
<i>Fomitiporia pentaphylacis</i>	Yuan 6012	China	JQ003900	JQ003901	Zhou and Xue (2012)
<i>Fomitiporia polymorpha</i>	MUCL 46166	USA	GU461955	DQ122393	Wu et al. (2022b)
<i>Fomitiporia polymorpha</i>	MUCL 46167	USA	GU461956	EF429233	Wu et al. (2022b)
<i>Fomitiporia pseudopunctata</i>	MUCL 38514	France	GU461953	AY618201	Wu et al. (2022b)
<i>Fomitiporia pseudopunctata</i>	MUCL 45670	France	GU461954	GU461980	Wu et al. (2022b)
<i>Fomitiporia pseudopunctata</i>	MUCL 51325	Czech	GU461948	GU461981	Wu et al. (2022b)
<i>Fomitiporia pseudopunctata</i>	MUCL 46168	France	JQ087891	JQ087918	Wu et al. (2022b)
<i>Fomitiporia punctata</i>	MUCL 47629	Japan	GU461950	GU461982	Amalfi et al. (2010)
<i>Fomitiporia punctata</i>	Dai 15772	China	KX639629	KX639647	Wu et al. (2022b)
<i>Fomitiporia punicata</i>	Cui 26	China	GU461975	GU461992	Wu et al. (2022b)
<i>Fomitiporia punicata</i>	Dai 7175	China	KX639632	KX639650	Wu et al. (2022b)
<i>Fomitiporia punicata</i>	Dai 10640	China	KX663825	KX639653	Wu et al. (2022b)
<i>Fomitiporia rhamnoides</i>	Dai 17369	China	MH234388	MH234391	Liu et al. (2018)
<i>Fomitiporia rhamnoides</i>	GL-2016	China	KT861405	KY697187	Liu et al. (2018)
<i>Fomitiporia rhamnoides</i>	Dai 18091	China	MH234389	MH234392	Liu et al. (2018)
<i>Fomitiporia robusta</i>	MUCL 51297	Estonia	JQ087892	JQ087919	Wu et al. (2022b)
<i>Fomitiporia robusta</i>	MUCL 51327	Czech	GU461949	GU461993	Wu et al. (2022b)
<i>Fomitiporia robustiformis</i>	JV 2402/8	Ecuador	PV368088	PV368094	Present study
<i>Fomitiporia robustiformis</i>	JV 2501/5-J	Peru	PV389821	—	Present study
<i>Fomitiporia roseo-bubalina</i>	JV 2402/1	Ecuador	PV368089	PV368095	Present study
<i>Fomitiporia roseo-bubalina</i>	JV 2402/10A	Ecuador	PV389820	PV389822	Present study
<i>Fomitiporia subtilissima</i>	FURB 47557	Brazil	NR_148098	NG_060426	GenBank
<i>Fomitiporia subtilissima</i>	FURB 47437	Brazil	KU557530	—	GenBank
<i>Fomitiporia sonora</i>	MUCL 47689	Arizona	JQ087893	JQ087920	Wu et al. (2022b)
<i>Fomitiporia</i> sp.	MUCL 53993	Mexico	JX093807	JX093851	Wu et al. (2022b)
<i>Fomitiporia</i> sp.	MUCL 53994	Mexico	JX093808	JX093852	Wu et al. (2022b)
<i>Fomitiporia</i> sp.	MUCL 53798	French Guiana	JX093811	JX093855	Wu et al. (2022b)
<i>Fomitiporia spinescens</i>	ICN 200566	Brazil	MN918549	—	Alves-Silva et al. (2020a)
<i>Fomitiporia subhippophaeicola</i>	Cui 12096	China	KU364421	KU364426	Chen et al. (2017)
<i>Fomitiporia subhippophaeicola</i>	Cui 12102	China	KU364423	KU364424	Chen et al. (2017)
<i>Fomitiporia subrobusta</i>	Dai 13576	China	KX639617	KX639635	Wu et al. (2022b)
<i>Fomitiporia subrobusta</i>	Dai 13577	China	KX639618	KX639636	Wu et al. (2022b)
<i>Fomitiporia subtropica</i>	Cui 9122	China	KX639622	KX639640	Wu et al. (2022b)
<i>Fomitiporia subtropica</i>	Cui 9115	China	KX639623	KX639641	Wu et al. (2022b)
<i>Fomitiporia tabaquilio</i>	MUCL 46230	Argentina	GU461940	DQ122394	Wu et al. (2022b)
<i>Fomitiporia tabaquilio</i>	MUCL 47754	Argentina	GU461941	GU461994	Wu et al. (2022b)
<i>Fomitiporia tasmanica</i>	Dai 18793	Australia	MH971173	MH971218	Wu et al. (2022b)

Species name	Sample no.	Location	GenBank accession no.		References
			ITS	nLSU	
<i>Fomitiporia tasmanica</i>	Dai 18799	Australia	MH971174	MH971219	Wu et al. (2022b)
<i>Fomitiporia tenuis</i>	MUCL 44802	Ethiopia	GU461957	AY618206	Amalfi et al. (2010)
<i>Fomitiporia tenuis</i>	MUCL 49948	Gabon	GU461958	GU461998	Amalfi et al. (2010)
<i>Fomitiporia tenuitubus</i>	Dai 16204	China	KX639619	KX639637	Wu et al. (2022b)
<i>Fomitiporia tenuitubus</i>	Yuan 5736	China	JQ003902	JQ003903	Wu et al. (2022b)
<i>Fomitiporia texana</i>	MUCL 47690	USA	JQ087894	JQ087921	Amalfi et al. (2012)
<i>Fomitiporia texana</i>	MUCL 51143	USA	JQ087895	JQ087922	Amalfi et al. (2012)
<i>Fomitiporia triqueter</i>	JV 2402/36	Ecuador	PV368090	—	Present study
<i>Fomitiporia triqueter</i>	JV 2402/68	Ecuador	PV368091	PV368096	Present study
<i>Fomitiporia tsitsikamensis</i>	CMW 47881	South Africa	MH599111	MH599121	Tchoumi et al. (2020)
<i>Fomitiporia tsitsikamensis</i>	CMW 48058	South Africa	MH599109	MH599120	Tchoumi et al. (2020)
<i>Fomitiporia tsitsikamensis</i>	CMW 48621	South Africa	MH599110	MH599123	Tchoumi et al. (2020)
<i>Fomitiporia tsugina</i>	MUCL 52702	USA	JQ087898	JQ087925	Amalfi et al. (2012)
<i>Fomitiporia tsugina</i>	MUCL 52703	USA	JQ087899	JQ087926	Amalfi et al. (2012)
<i>Fomitiporia uncinata</i>	ICN 200561	Brazil	—	MN918542	Alves-Silva et al. (2020a)
<i>Neophellinus uncisetus</i>	MUCL 46231	Argentina	GU461960	EF429235	Wu et al. (2022b)
<i>Neophellinus uncisetus</i>	MUCL 47061	Argentina	GU461972	GU462000	Wu et al. (2022b)

Bold = new taxa; — refers to the data unavailability.

chains and starting from random trees. Trees were sampled every 100 generations from a total of 6 million generations. The first 25% of trees were removed, and the remaining ones were used to construct a 50% majority consensus tree and calculate Bayesian posterior probabilities (BPPs). All trees were viewed in FigTree 1.4.3. The two methods constructed nearly congruent topologies for each alignment. Therefore, only the topologies inferred from the ML method were presented along with bootstrap values from the ML method ($\geq 50\%$) and BPPs (≥ 0.9) from the BI methods at the nodes.

Results

Molecular phylogeny

The combined ITS and nLSU dataset included sequences from 123 fungal collections representing 65 taxa of *Fomitiporia*, and two samples of the genus *Neophellinus* were used as the outgroup. The final alignment comprised a total of 1,827 base pairs (bp), including 930 bp of ITS and 898 bp of nLSU. The best model was estimated and applied in the Bayesian analysis: GTR+I+G, lset nst = 6, rates = invgamma; prset statefreqpr = dirichlet (1,1,1,1). Bayesian analysis yielded a nearly congruent topology (average standard deviation of split frequencies = 0.009985) with the ML analysis; therefore, only the topology from the ML analysis is presented. The current phylogeny (Fig. 1) showed that the newly sequenced specimens fell into the *Fomitiporia* clade and formed three independent new lineages with strong support: *Fomitiporia robustiformis* (BS = 99% in ML, BPP = 1.00), *Fomitiporia roseo-bubalina* (BS = 100% in ML, BPP = 1.00), and *Fomitiporia triqueter* (BS = 100% in ML, BPP = 1.00).

The ITS and nLSU sequences generated in this study were subjected to BLAST searches in the NCBI database (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). The BLASTN results are consistent with our phylogeny, and the top five search results for the type specimen sequences of the three new *Fomitiporia* species are provided (Tables 2, 3).



Figure 1. Maximum likelihood tree illustrating the phylogeny of *Fomitiporia* based on the combined dataset of ITS + nLSU sequences. The maximum likelihood bootstrap values (≥ 50) and Bayesian posterior probability values (≥ 0.90) are indicated above the branches. The new species is in bold.

Table 2. BLAST results in the NCBI database with ITS.

Species	Match rank	Matched species	Identity (%)	Score	Query coverage (%)	Alignment length	GenBank accession no.	Origin
<i>Fomitiporia robustiformis</i> (holotype: JV 2402/8)	1	<i>Phellinus texanus</i>	93.57	1123	100	783	JQ087895	USA
	2	<i>Phellinus texanus</i>	93.57	1122	100	783	JQ087894	USA
	3	<i>Fomitiporia deserticola</i>	93.83	1110	98	744	KT381634	USA
	4	<i>Fomitiporia deserticola</i>	93.83	1109	98	743	KT381635	USA
	5	<i>Phellinus texanus</i>	93.73	1105	98	749	KT381636	USA
<i>Fomitiporia roseo-bubalina</i> (holotype: JV 2402/1)	1	<i>Fomitiporia mediterranea</i>	86.35	821	100	767	PQ555869	—
	2	<i>Fomitiporia mediterranea</i>	86.30	817	100	801	AY849303	Italy
	3	<i>Phellinus erectus</i>	86.30	817	100	778	GU461939	France
	4	<i>Fomitiporia mediterranea</i>	86.22	815	100	781	EF442330	—
	5	<i>Fomitiporia mediterranea</i>	86.22	815	100	766	PQ060508	Italy
<i>Fomitiporia triqueter</i> (holotype: JV 2402/68)	1	<i>Fomitiporia baccharidis</i>	93.38	1085	99	738	MW880690	—
	2	<i>Phellinus tabaquilio</i>	92.82	1070	99	762	GU461940	Argentina
	3	<i>Phellinus tabaquilio</i>	92.80	1064	99	756	GU461941	Argentina
	4	<i>Fomitiporia apiahyna</i>	90.31	976	99	770	MN918572	Brazil
	5	<i>Fomitiporia nubicola</i>	91.09	972	96	735	KU663308	Brazil

Table 3. BLAST results in the NCBI database with nLSU.

Species	Match rank	Matched species	Identity (%)	Score	Query coverage (%)	Alignment length	GenBank accession no.	Origin
<i>Fomitiporia robustiformis</i> (holotype: JV 2402/8)	1	<i>Phellinus hartigii</i>	97.58	1620	99	1459	PV336067	Poland
	2	<i>Fomitiporia tsugina</i>	97.58	1620	99	978	KC551838	—
	3	<i>Phellinus hartigii</i>	97.58	1620	99	978	KC551856	—
	4	<i>Phellinus robustus</i>	97.58	1620	99	1459	PV250053	Poland
	5	<i>Fomitiporia punctata</i>	97.58	1620	99	1459	PV271020	Poland
<i>Fomitiporia roseo-bubalina</i> (holotype: JV 2402/1)	1	<i>Fomitiporia punctata</i>	97.87	1615	99	1459	PV271020	Poland
	2	<i>Fomitiporia tsugina</i>	97.76	1609	99	978	KC551836	—
	3	<i>Fomitiporia tsugina</i>	97.76	1609	99	978	KC551838	—
	4	<i>Fomitiporia punctata</i>	97.76	1609	99	1459	PV277016	Poland
	5	<i>Phellinus robustus</i>	97.76	1609	99	1459	PV250053	Poland
<i>Fomitiporia triqueter</i> (holotype: JV 2402/68)	1	<i>Fomitiporia baccharidis</i>	98.32	1663	100	957	MW880691	—
	2	<i>Fomitiporia tsugina</i>	98.41	1661	99	978	KC551836	—
	3	<i>Fomitiporia tsugina</i>	98.31	1657	99	978	KC551844	—
	4	<i>Fomitiporia tsugina</i>	98.31	1657	99	978	KC551839	—
	5	<i>Fomitiporia tsugina</i>	98.31	1657	99	978	KC551846	—

Taxonomy

***Fomitiporia robustiformis* Jian Chen, Yuan Yuan, K.Y. Luo, Y.C. Dai & Vlasák, sp. nov.**
MycoBank No: 858483
Figs 2, 3

Diagnosis. *Fomitiporia robustiformis* is closely related to *F. texana* (Murrill) Nuss and *F. deserticola* Vlasák, but *F. texana* differs from *F. robustiformis* by the presence of hymenial setae and cystidioles, larger basidiospores (7–9 × 6.5–9 µm vs. 5–6.4 × 4.7–6.1 µm). *F. deserticola* is distinguished from *F. robustiformis* by the presence of abundant cystidioles and larger basidiospores (6–7.5 × 5.5–7 µm vs. 5–6.4 × 4.7–6.1 µm).
Holotype. ECUADOR • Arenillas, dry tropical forest, on living *Acacia* sp., 17.II.2024, leg J. Vlasák, JV 2402/8 (BJFC 053711).
Etymology. *Robustiformis* (Lat.) refers to a new species characterized by robust basidiomata.



Figure 2. Basidiomata of *Fomitiporia robustiformis* (holotype, JV 2402/8). Scale bar: 1 cm.

Description. Basidiomata perennial, pileate, inseparable, sessile, without distinctive odor or taste when fresh, woody hard in consistency, light in weight when dry; pilei ungulate, fan-shaped, projecting up to 5 cm, 5 cm wide, and 3 cm thick at base; pileal surface brown, sometimes greenish from algae, glabrous; margin broad and rounded. Pore surface pale mouse-gray when fresh, becoming yellowish brown when dry, not glossy; sterile margin brown, up to 1 mm wide; pores circular to angular, 6–7 per mm; dissepiments thin, entire. Context yellow-brown, woody hard, up to 7 mm thick; tubes pale gray to yellow-brown, paler than pore surface, woody hard, up to 2 cm long, annual layers indistinct.

Hyphal structure. Hyphal system dimitic, generative hyphae simple septate; all hyphae IKI–, CB–; tissue becoming dark brown in KOH.

Context. Generative hyphae frequent, pale yellow, slightly thick-walled, rarely branched, frequently septate, 3–3.5 µm in diam; skeletal hyphae dominant, yellowish brown, thick-walled, unbranched, more or less flexuous, interwoven, 3–4.5 µm in diam.

Tubes. Generative hyphae pale yellowish, slightly thick-walled, rarely branched, frequently septate, 3–3.5 µm in diam; skeletal hyphae dominant, yellowish, thick-walled, unbranched, more or less straight, parallel along the tubes, 3–4.5 µm in diam. Hymenial setae absent; cystidioles present, fusoid, hyaline, thin-walled, 14–18 × 2.5–3.5 µm; basidia clavate, with four sterigmata and a simple septum at the base, 10–18 × 6–7.5 µm; basidioles dominant in hymenium, in shape similar to basidia, but slightly smaller.

Basidiospores globose, hyaline, thick-walled, smooth, some with a guttule, IKI[+], CB+, (4.4–)5–6.4(–7.1) × (4.4–)4.7–6.1(–6.9) µm, L = 5.66 µm, W = 5.39 µm, Q = 1.05 (n = 30/1).




Figure 3. Microscopic structures of *Fomitiporia robustiformis* (Holotype, JV 2402/8) **a** basidiospores **b** hyphae from context **c** basidia and basidioles **d** cystidioles **e** hyphae from the tube trama.

Type of rot. White rot.

Additional specimen (paratype) examined. PERU • Chongoyape, dry tropical forest, on a dead branch of a living angiosperm tree, 21.I.2025, leg J. Vlasák Jr., JV 2501/5-J.

***Fomitiporia roseo-bubalina* Jian Chen, Yuan Yuan, K.Y. Luo, Y.C. Dai & Vlasák, sp. nov.**
MycoBank No: 858484
Figs 4, 5

Diagnosis. *Fomitiporia roseo-bubalina* is related to *F. ovoidospora* Y.C. Dai & F. Wu, but *F. ovoidospora* differs from *F. roseo-bubalina* by perennial basidiocarps,



Figure 4. Basidiomata of *Fomitiporia roseo-bubalina* (Holotype, JV 2402/1). Scale bar: 1 cm.

the presence of fusoid cystidioles, smaller pores (9–10 per mm vs. 4–5 per mm), and smaller basidiospores ($4.7\text{--}5.5 \times 3.8\text{--}5 \mu\text{m}$ vs. $5.3\text{--}6.7 \times 5\text{--}6.2 \mu\text{m}$).

Holotype. ECUADOR • Arenillas, dry tropical forest, on the base of living *Acacia*, 17.II.2024, leg J. Vlasák, JV 2402/1 (BJFC 053710).

Etymology. *Roseo-bubalina* (Lat.) refers to the new species characterized by pink-buff pores when dry.

Description. Basidiomata annual to perennial, resupinate to pileate, inseparable, growing around bark fissures at the base of living trees in shelf-like, vertical assemblages of triquetrous pilei, without distinctive odor or taste when fresh, woody hard in consistency; pilei slightly ungulate, projecting up to 2.5 cm, 3 cm wide, and 0.5 cm thick at base; pileal surface grayish, glabrous; margin blunt. Pore surface pink buff when dry, glossy; sterile margin cream, up to 1 mm wide; pores circular to angular, 4–5 per mm; dissepiments thin, entire. Context very thin; tubes yellowish brown, paler than pore surface, woody hard, up to 2 mm long, annual layers indistinct.

Hyphal structure. Hyphal system dimitic, generative hyphae simple septate; all hyphae IKI–, CB–; tissue becoming dark brown in KOH.

Tubes. Generative hyphae pale yellowish, slightly thick-walled, rarely branched, frequently septate, $2.5\text{--}3.5 \mu\text{m}$ in diam; skeletal hyphae dominant, yellowish, thick-walled, unbranched, more or less straight, parallel along the tubes, $3.5\text{--}4.8 \mu\text{m}$ in diam. Hymenial setae absent; cystidia and cystidioles absent; basidia subglobose to clavate, with four sterigmata and a simple septum at the base, $10\text{--}18 \times 6\text{--}9 \mu\text{m}$; basidioles dominant in hymenium, in shape similar to basidia, but slightly smaller.




Figure 5. Microscopic structures of *Fomitiporia roseo-bubalina* (Holotype, JV 2402/1) **a** basidiospores **b** basidia and basidioles **c** hyphae from the tube trama.

Basidiospores globose, hyaline, thick-walled, smooth, IKI[+], CB+, (5–)5.3–6.7(–7.4) × (4.9–)5–6.2(–6.6) µm, L = 6.01 µm, W = 5.60 µm, Q = 1.07 (n = 30/1).

Type of rot. White rot.

Additional specimen (paratype) examined. ECUADOR • Arenillas, dry tropical forest, on the base of living *Acacia*, 17.II.2024, leg J. Vlasák, JV 2402/10A.

***Fomitiporia triqueter* Jian Chen, Yuan Yuan, K.Y. Luo, Y.C. Dai & Vlasák, sp. nov.**

MycoBank No: 858485

Figs 6, 7

Diagnosis. *Fomitiporia triqueter* is closely related to *F. baccharidis* and *F. tabaquilio* (Urcelay, Robledo & Rajchenb.) Decock & Robledo, and they all occur in South America. However, the latter two species differ from *F. triqueter* in having larger basidiospores (5.5–6.5 × 4.8–5.8 µm in *F. baccharidis*, 6–7.5 × 8–6.5 µm in *F. tabaquilio*, vs. 3.4–5.5 × 3.2–5.4 µm in *F. triqueter*).

Holotype. ECUADOR • Macará, dry tropical forest, on the thick bark of a living angiosperm tree, 23.II.2024, leg J. Vlasák Jr., JV 2402/68 (BJFC 053713).

Etymology. *Triqueter* (Lat.) refers to the new species characterized by triqueter basidiomata in section.

Description. Basidiomata biennial to perennial, but evidently short-living, pileate, without distinctive odor or taste when fresh, woody hard in consistency; pilei ungulate, triquetrous in section, projecting up to 4 cm, 2.5 cm wide, and 2 cm thick at base; pileal surface grayish dark, concentrically sulcate,



Figure 6. Basidiomata of *Fomitiporia triqueter* (holotype, JV 2402/68). Scale bar: 1 cm.




Figure 7. Microscopic structures of *Fomitiporia triqueter* (holotype, JV 2402/68) **a** basidiospores **b** basidia and basidioles **c** hyphae from context **d** hyphae from the tube trama.

glabrous, soon cracked and blackish with age; margin sharp; pore surface yellowish brown when fresh, becoming brown when dry; sterile margin yellowish brown, up to 100 µm wide; pores circular, 9–10 per mm; dissepiments slightly thick, entire. Context yellowish brown, woody hard, up to 2 mm thick; tubes yellowish brown, paler than pore surface, woody hard, up to 2 cm long, annual layers indistinct.

Hyphal structure. Hyphal system dimitic, generative hyphae simple septate; all hyphae IKI–, CB–; tissue becoming dark brown in KOH.

Context. Generative hyphae frequent, pale yellow, slightly thick-walled, occasionally branched, frequently septate, 2.5–3 µm in diam; skeletal hyphae

dominant, yellowish brown, thick-walled, unbranched, more or less flexuous, interwoven, 3–4 µm in diam.

Tubes. Generative hyphae golden, slightly thick-walled, occasionally branched, frequently septate, 2.5–3.5 µm in diam; skeletal hyphae dominant, yellowish, thick-walled with a medium lumen, unbranched, more or less straight, parallel along the tubes, 3–4.5 µm in diam. Hymenial setae absent; cystidia and cystidioles absent; basidia subglobose to capitate, with four sterigmata and a simple septum at the base, 13–18 × 7–11 µm; basidioles dominant in hymenium, in shape similar to basidia, but slightly smaller; large rhomboid crystals present in the hymenium.

Basidiospores globose, hyaline, thick-walled, smooth, IKI[+], slightly CB+, (3.1–)3.4–5.5(–5.7) × (2.9–)3.2–5.4(–5.8) µm, L = 4.57 µm, W = 4.39 µm, Q = 1.04–1.05 (n = 60/2).

Type of rot. White rot in the bark; no signs of wood rot under the infested bark.

Additional specimen (paratype) examined. ECUADOR • Macará, dry tropical forest, on the thick bark of a living angiosperm tree, 21.II.2024, leg J. Vlasák Jr., JV 2402/36 (BJFC 053712).

Discussion

Four species of *Fomitiporia* from Ecuador have been reported in the last 10 years, including *F. baccharidis* (Pat.) Decock et al., *F. conyana* Alves-Silva & Drechsler-Santos, *F. nubicola* Alves-Silva et al., and *F. impercepta* Morera et al. (Wu et al. 2022b). In the present study, three new species of *Fomitiporia* are described, all of which share the typical characteristics of *Fomitiporia*, including subglobose, hyaline, thick-walled, smooth, dextrinoid, and cyanophilous basidiospores. The phylogenetic analysis revealed that they are distantly related to other taxa in the genus (Fig. 1).

Phylogenetically, *Fomitiporia robustiformis* is closely related to *F. texana* (Murrill) Nuss and *F. deserticola* Vlasák (Fig. 1), but *F. texana* differs from *F. robustiformis* by the presence of hymenial setae and cystidioles, larger pores (4–6 per mm vs. 6–7 per mm), larger subglobose basidiospores (7–9 × 6.5–9 µm vs. 5–6.4 × 4.7–6.1 µm), and occurrence in the Southwest USA (Larsen and Cobb-Pouille 1990). *F. deserticola* is distinguished from *F. robustiformis* by the presence of rare hymenial setae and abundant cystidioles, larger pores (4–6 per mm vs. 6–7 per mm), larger and subglobose basidiospores (6–7.5 × 5.5–7 µm vs. 5–6.4 × 4.7–6.1 µm), and a distribution in Arizona, USA (Vlasák and Vlasák 2016). Morphologically, *F. robustiformis* is most similar to *F. subrobusta* B.K. Cui & Hong Chen by sharing perennial, pileate basidiocarps; approximately the same size pores; the presence of cystidioles; and the absence of hymenial setae. However, *F. subrobusta* has larger and subglobose to obovoid basidiospores (6.2–6.8 × 5.2–6 µm), and its distribution is in South China (Chen and Cui 2017).

Morphologically, *Fomitiporia roseo-bubalina* is characterized by annual to perennial basidiocarps, absent hymenial setae, and globose basidiospores, which is similar to *F. pentaphylacis* L.W. Zhou (Zhou and Xue 2012). However, *F. pentaphylacis* differs from *F. roseo-bubalina* by its smaller pores (6–9 per mm vs. 4–5 per mm), larger basidiospores (5.9–7.6 × 5.4–6.5 µm vs. 5.3–6.7 × 5–6.2 µm), the presence of cystidioles, and its distribution in South China. Phylogenetically, *F. roseo-bubalina* is related to *F. ovoidospora* Y.C. Dai & F. Wu, but *F. ovoidospora* differs from *F. roseo-bubalina* by perennial basidiocarps, dimidiate or triquetrous and cracked pilei, the presence of fusoid cystidioles, smaller

pores (9–10 per mm vs. 4–5 per mm), and smaller and ovoid basidiospores ($4.7\text{--}5.5 \times 3.8\text{--}5 \mu\text{m}$ vs. $5.3\text{--}6.7 \times 5\text{--}6.2 \mu\text{m}$). Furthermore, *F. ovoidospora* grows on *Khaya* or other angiosperm wood and has a distribution in Vietnam.

Fomitiporia triqueter is characterized by biennial to perennial and pileate basidiomata, ungulate to triquetrous and cracked pileal, the absence of hymenial setae and cystidioles, the presence of large rhomboid crystals in the hymenium, and small and globose basidiospores ($3.4\text{--}5.5 \times 3.2\text{--}5.4 \mu\text{m}$). *F. triqueter* and *F. ovoidospora* share similar pores and basidiospores, but the latter has cystidioles and lacks rhomboid crystals (Wu et al. 2022b). Phylogenetically, *F. triqueter* is closely related to *F. baccharidis* and *F. tabaquilio* (Urcelay, Robledo & Rajchenb) Decock & Robledo, and they all occur in South America. However, *F. baccharidis* is different from *F. triqueter* by its velutinate and uncracked pileal, strongly glossy pore surface, and larger basidiospores ($5.5\text{--}6.5 \times 4.8\text{--}5.8 \mu\text{m}$ vs. $3.4\text{--}5.5 \times 3.2\text{--}5.4 \mu\text{m}$, Amalfi et al. 2014). *Fomitiporia tabaquilio* differs from *F. triqueter* by its glossy pore surface, the presence of cystidioles, larger and subglobose basidiospores ($6\text{--}7.5 \times 8\text{--}6.5 \mu\text{m}$ vs. $3.4\text{--}5.5 \times 3.2\text{--}5.4 \mu\text{m}$), and its distribution in Argentina (Decock et al. 2007).

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The research is supported by the Young Elite Scientists Sponsorship Program by CAST (No. 2023QNRC001). The research of J. Vlasák was financed by the institutional support of the Academy of Sciences of the Czech Republic (RVO: 60077344).

Author contributions

All authors designed the research and contributed to data analysis and interpretation, and prepared the samples and drafted the manuscript.

Author ORCIDs

Jian Chen  <https://orcid.org/0000-0001-8409-7791>

Josef Vlasák  <https://orcid.org/0000-0002-5363-4459>

Kai-Yue Luo  <https://orcid.org/0000-0001-6145-4864>

Yuan Yuan  <https://orcid.org/0000-0001-6674-9848>

Data availability

All of the data that support the findings of this study are available in the main text.

References

Alves-Silva G, Drechsler-Santos ER, Da Silveira RMB (2020a) Bambusicolous *Fomitiporia* revisited: Multilocus phylogeny reveals a clade of host-exclusive species. *Mycologia* 112(3): 633–648. <https://doi.org/10.1080/00275514.2020.1741316>

- Alves-Silva G, Reck MA, Da Silveira RMB, Bittencourt F, Robledo GL, Góes-Neto A, Drechsler-Santos ER (2020b) The neotropical *Fomitiporia* (Hymenochaetales, Basidiomycota): The redefinition of *F. apiahyna* s.s. allows revealing a high hidden species diversity. *Mycological Progress* 19(8): 769–790. <https://doi.org/10.1007/s11557-020-01593-5>
- Amalfi M, Decock C (2014) *Fomitiporia expansa*, an undescribed species from French Guiana. *Cryptogamie. Mycologie* 35(1): 73–85. <https://doi.org/10.7872/crym.v35.iss1.2014.73>
- Amalfi M, Yombiyeni P, Decock C (2010) *Fomitiporia* in sub-Saharan Africa: Morphology and multigene phylogenetic analysis support three new species from the Guineo-Congolian rainforest. *Mycologia* 102(6): 1303–1317. <https://doi.org/10.3852/09-083>
- Amalfi M, Raymundo T, Valenzuela R, Decock C (2012) *Fomitiporia cupressicola* sp. nov., a parasite on *Cupressus arizonica*, and additional unnamed clades in the southern USA and northern Mexico, determined by multilocus phylogenetic analyses. *Mycologia* 104(4): 880–893. <https://doi.org/10.3852/11-196>
- Amalfi M, Robledo G, Decock C (2014) *Fomitiporia baccharidis* comb. nov., a little known species from high elevation Andean forests and its affinities within the neotropical *Fomitiporia* lineages. *Mycological Progress* 13(4): 995. <https://doi.org/10.1007/s11557-014-0995-x>
- Brown AA, Lawrence DP, Baumgartner K (2020) Role of basidiomycete fungi in the grapevine trunk disease esca. *Plant Pathology* 69(2): 205–220. <https://doi.org/10.1111/ppa.13116>
- Chen H, Cui BK (2017) Multi-locus phylogeny and morphology reveal five new species of *Fomitiporia* (Hymenochaetaceae) from China. *Mycological Progress* 16(7): 687–701. <https://doi.org/10.1007/s11557-017-1306-0>
- Chen JJ, Cui BK, Zhou LW, Korhonen K, Dai YC (2015) Phylogeny, divergence time estimation, and biogeography of the genus *Heterobasidion* (Basidiomycota, Russulales). *Fungal Diversity* 71: 185–200. <https://doi.org/10.1007/s13225-014-0317-2>
- Chen H, Zhou JL, Cui BK (2017) Two new species of *Fomitiporia* (Hymenochaetales, Basidiomycota) from Tibet, southwest China. *Mycologia* 108(5): 1010–1017. <https://doi.org/10.3852/16-011>
- Chen JJ, Wu YD, Ji XH, Gates G, Xu X (2021) A new species of *Fomitiporia* (Hymenochaetales) from Australia. *Phytotaxa* 489(2): 200–208. <https://doi.org/10.11646/phytotaxa.489.2.8>
- Cloete M, Fischer M, Mostert L, Halleen F (2014) A novel *Fomitiporia* species associated with esca on grapevine in South Africa. *Mycological Progress* 13(2): 303–311. <https://doi.org/10.1007/s11557-013-0915-5>
- Dai YC (2022) New trend of edible and medicinal fungi research in China—based on publications in *Mycosystema* during 2000–2021. *Journal of Fungal Research* 20: 141–156. <https://doi.org/10.13341/j.jfr.2022.1494>
- Dai YC (2023) Research progress on polypore domesticated cultivation in China. *Journal of Fungal Research* 21: 151–156. <https://doi.org/10.13341/j.jfr.2023.1602>
- Dai YC, Cui BK (2011) *Fomitiporia ellipsoidea* has the largest fruiting body among the fungi. *Fungal Biology* 115: 813–814. <https://doi.org/10.1016/j.funbio.2011.06.008>
- Dai YC, Cui BK, Yuan HS, Li BD (2007) Pathogenic wood-decaying fungi in China. *Forest Pathology* 37(2): 105–120. <https://doi.org/10.1111/j.1439-0329.2007.00485.x>
- Dai YC, Yang ZL, Cui BK, Yu CJ, Zhou LW (2009) Species diversity and utilization of medicinal mushrooms and fungi in China. *International Journal of Medicinal Mushrooms* 11: 287–302. [review] <https://doi.org/10.1615/IntJMedMushr.v11.i3.80>

- Dai YC, Yang ZL, Cui BK, Wu G, Yuan HS, Zhou LW, He SH, Ge ZW, Wu F, Wei YL, Yuan Y, Si J (2021) Diversity and systematics of the important macrofungi in Chinese forests. *Mycosystema* 40: 770–805. <https://doi.org/10.13346/j.mycosystema.210036>
- Decock C, Figueroa SH, Robledo G, Castillo G (2007) *Fomitiporia punctata* (Basidiomycota, Hymenochaetales) and its presumed taxonomic synonyms in America: Taxonomy and phylogeny of some species from tropical/subtropical areas. *Mycologia* 99(5): 733–752. <https://doi.org/10.1080/15572536.2007.11832537>
- Dong JH, Li Q, Yuan Q, Luo YX, Zhang XC, Dai YF, Zhou Q, Liu XF, Deng YL, Zhou HM, Muhammad A, Zhao CL (2024) Species diversity, taxonomy, molecular systematics and divergence time of wood-inhabiting fungi in Yunnan-Guizhou Plateau, Asia. *Mycosphere* 15(1): 1110–1293. <https://doi.org/10.5943/mycosphere/15/1/10>
- Du R, Wu F, Gate GM, Dai YC, Tian XM (2020) Taxonomy and phylogeny of *Sidera* (Hymenochaetales, Basidiomycota): Four new species and keys to species of the genus. *MycoKeys* 68: 115–135. <https://doi.org/10.3897/mycokeys.68.53561>
- Fischer M (1996) On the species complexes within *Phellinus*: *Fomitiporia* revisited. *Mycological Research* 100(12): 1459–1467. [https://doi.org/10.1016/S0953-7562\(96\)80078-5](https://doi.org/10.1016/S0953-7562(96)80078-5)
- Fischer M (2002) A new wood-decaying basidiomycete species associated with esca of grapevine: *Fomitiporia mediterranea* (Hymenochaetales). *Mycological Progress* 1(3): 315–324. <https://doi.org/10.1007/s11557-006-0029-4>
- Fischer M, Binder M (2004) Species recognition, geographic distribution and host-pathogen relationships: A case study in a group of lignicolous basidiomycetes, *Phellinus* s.l. *Mycologia* 96(4): 799–811. <https://doi.org/10.1080/15572536.2005.11832927>
- Ghobad-Nejhad M, Zhou LW, Tomšovský M, Angelini P, Cusumano G, Angeles Flores G, Venanzoni R, Wang XW, Chaharmiri-Dokhaharani S, Moridi Farimani M, Pärtel K, Dai YC, Wu F (2024) Unlocking nature's pharmacy: Diversity of medicinal properties and mycochemicals in the family Hymenochaetaceae (Agaricomycetes, Basidiomycota). *Mycosphere* 15(1): 6347–6438. <https://doi.org/10.5943/mycosphere/15/1/27>
- Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology* 52(5): 696–704. <https://doi.org/10.1080/10635150390235520>
- Hall TA (1999) Bioedit: A user-friendly biological sequence alignment editor and analysis program for windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98. <https://doi.org/10.1021/bk-1999-0734.ch008> [oxford] <https://doi.org/10.1021/bk-1999-0734.ch008>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Larsen MJ, Cobb-Poullé LA (1990) *Phellinus* (Hymenochaetaceae). A survey of the world taxa. *Mycologia* 82(6): 801. <https://doi.org/10.2307/3760175>
- Li GJ, Hyde KD, Zhao RL, Hongsan S, Abdel-Aziz FA, Abdel-Wahab MA, Alvarado P, Alves-Silva G, Ammirati JF, Ariyawansa HA, Baghela A, Bahkali AH, Beug M, Bhat DJ, Bojantchev D, Boonpratuang T, Bulgakov TS, Camporesi E, Boro MC, Ceska O, Chakraborty D, Chen JJ, Chethana KWT, Chomnunti P, Consiglio G, Cui BK, Dai DQ, Dai YC, Daranagama DA, Das K, Dayarathne MC, De Crop E, De Oliveira RJV, de Souza CAF, de Souza JI, Dentinger BTM, Dissanayake AJ, Doilom M, Drechsler-Santos ER, Ghobad-Nejhad M, Gilmore SP, Góes-Neto A, Gorczak M, Haitjema CH, Hapuarachchi KK, Hashimoto A, He MQ, Henske JK, Hirayama K, Iribarren MJ, Jayasiri SC, Jayawardena RS, Jeon SJ, Jerônimo GH, Jesus AL, Jones EBG, Kang JC, Karunarathna SC, Kirk PM, Konta S, Kuhnert E, Langer E, Lee HS, Lee HB, Li WJ, Li XH, Liimatainen K, Lima DX,

- Lin CG, Liu JK, Liu XZ, Liu ZY, Luangsa-Ard JJ, Lücking R, Lumbsch HT, Lumyong S, Leaño EM, Marano AV, Matsumura M, Mckenzie EHC, Mongkolsamrit S, Mortimer PE, Nguyen TTT, Niskanen T, Norphanphoun C, O'Malley MA, Parnmen S, Pawłowska J, Perera RH, Phookamsak R, Phukhamsakda C, Pires-Zottarelli CLA, Raspé O, Reck MA, Rocha SCO, de Santiago ALCM, Senanayake IC, Setti L, Shang QJ, Singh SK, Sir EB, Solomon KV, Song J, Srikitikulchai P, Stadler M, Suetrong S, Takahashi H, Takahashi T, Tanaka K, Tang LP, Thambugala KM, Thanakitpipattana D, Theodorou MK, Thongbai B, Thummarukcharoen T, Tian Q, Tibpromma S, Verbeken A, Vizzini A, Vlasák J, Voigt K, Wanasinghe DN, Wang Y, Weerakoon G, Wen HA, Wen TC, Wijayawardene NN, Wongkanoun S, Wrzosek M, Xiao YP, Xu JC, Yan JY, Yang J, Da Yang S, Hu Y, Zhang JF, Zhao J, Zhou LW, Peršoh D, Phillips AJL, Maharachchikumbura SSN (2016) Fungal diversity notes 253–366: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 78(1): 1–237. <https://doi.org/10.1007/s13225-016-0366-9>
- Liu TZ, Chen Q, Han ML, Wu F (2018) *Fomitiporia rhamnoides* sp. nov. (Hymenochaetales, Basidiomycota), a new polypore growing on *Hippophae* from China. *MycKeys* 36: 35–43. <https://doi.org/10.3897/mycokeys.36.25986>
- Liu ZB, Yuan Y, Dai YC, Liu HG, Vlasák J, Zeng GY, He SH, Wu F (2025) Global diversity and systematics of Hymenochaetaceae with non-poroid hymenophore. *Fungal Diversity* 131: 1–97. <https://doi.org/10.1007/s13225-025-00552-3>
- Maddison WP, Maddison DR (2017) Mesquite: A modular system for evolutionary analysis, Version 3.2. <http://mesquiteproject.org>
- Manawasinghe IS, Hyde KD, Wanasinghe DN, Karunarathna SC, Maharachchikumbura SSN, Samarakoon MC, Voglmayr H, Pang K, Chiang MW, Jones EBG, Saxena RK, Kumar A, Rajeshkumar KC, Selbmann L, Coleine C, Hu Y, Ainsworth AM, Liimatainen K, Niskanen T, Ralaiveloarisoa A, Arumugam E, Kezo K, Kaliyaperumal M, Gunaseelan S, Dissanayake AJ, Khalid AN, Gajanayake AJ, Flakus A, Armand A, Aptroot A, Rodrigues A, Tsurukau A, López-Villalba Á, de Farias ARG, Sánchez A, Góes-Neto A, Goto BT, de Souza CAF, Chuaseeharonnachai C, Lin C, Li C, Denchev CM, Guerra-Mateo D, Tennakoon DS, Wei D, Begerow D, Alves E, Drechsler-Santos ER, Sousa ES, de Medeiros EV, Langer E, Zhang F, de Souza FA, Magurno F, Barreto GG, Moreno G, Mane G, Alves-Silva G, Da Silva GA, Xia G, Shen H, Gui H, Senanayake IC, Luangsa-Ard JJ, Liu J, Liu J, Ma J, Lin J, Beserra Jr JEA, Cano-Lira JF, Gené J, Harikrishnan K, Lu L, Dos Santos LA, Xu L, Lacerda LT, Gusmão LFP, Cáceres MES, Câmara MPS, de Barros-Barreto MBB, Calabon MS, Kukwa M, Kemler M, de Melo MP, Ghobad-Nejhad M, Luo M, Ding M, Doilom M, Phonemany M, Usman M, Thongklang N, Boonyuen N, Ashtekar N, Kulathnaga ND, Sruthi OP, Kwantong P, Ansil PA, Kooij PW, Zhao Q, Alfenas RF, de Oliveira RJV, Singh R, Da Silva RMF, Avchar R, Morey R, Sharma R, Xu R, Da Silveira RMB, Xu R, Jayawardena RS, Nanu S, Nuankaew S, Tibpromma S, Boonmie S, Somrithipol S, Varghese S, Moreira SI, Rajwar S, He S, Kumar TKA, Denchev TT, Luangharn T, de Oliveira TGL, Du T, Wen T, Du T, Wu T, Sri-Indrasutdhi V, Doyle VP, Baulin V, Dong W, Li W, Lu W, Tian W, Dos Vieira WA, von Brackel W, Yu X, Zhang X, Liu X, Peng X, Chen Y, Yang Y, Gao Y, Xiong Y, Shu Y, Lu Y, Shen Y, Zhou Y, Zhang YX, Zhang W, Luo Z, Madushani MA, Cheewangkoon R, Song JG, Xu B (2024) Fungal diversity notes 1818–1918: Taxonomic and phylogenetic contributions on genera and species of fungi. *Fungal Diversity* 130: 1–261. <https://doi.org/10.1007/s13225-024-00541-y>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>

- Murrill WA (1907) Polyporaceae. North American Flora 9: 1–131.
- Petersen J (1996) The Danish Mycological Society's colour-chart. Foreningen til Svampekundskabens Fremme, Greve.
- Posada D (2008) jModelTest: Phylogenetic model averaging. Molecular Biology and Evolution 25(7): 1253–1256. <https://doi.org/10.1093/molbev/msn083>
- Rajchenberg M, Pildain MB, Madriaga DC, de Errasti A, Riquelme C, Becerra J (2019) New Poroid Hymenochaetaceae (Basidiomycota, Hymenochaetales) from Chile. Mycological Progress 18(6): 865–877. <https://doi.org/10.1007/s11557-019-01495-1>
- Shen RM, Lu JY, Zhang Y, Wei H, Lu L (2024) Hypolipidemic functions of edible and medicinal mushrooms: A review. Mycosystema 43(8): 240050. <https://doi.org/10.13346/j.mycosystema.240050>
- Tchoumi JMT, Coetzee MPA, Rajchenberg M, Roux J (2020) Poroid Hymenochaetaceae associated with trees showing wood-rot symptoms in the Garden Route National Park of South Africa. Mycologia 112(4): 722–741. <https://doi.org/10.1080/00275514.2020.1753160>
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The Clustal_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Research 25(24): 4876–4882. <https://doi.org/10.1093/nar/25.24.4876>
- Viner I (2025) Taxonomy and phylogeny of Hyphodontia sensu lato (Hymenochaetales, Basidiomycota). Towards the family level classification of Hymenochaetales. University of Helsinki, Dissertationes Universitatis Helsingiensis 61/2025. <http://hdl.handle.net/10138/591580>
- Vlasák J, Vlasák J (2016) Two new polypore species from the southwestern USA: *Fomitiporia fissurata* and *F. deserticola*. Mycotaxon 131(1): 193–203. <https://doi.org/10.5248/131.193>
- Wagner T, Fischer M (2001) Natural groups and a revised system for the European poroid Hymenochaetales (Basidiomycota) supported by nLSU rDNA sequence data. Mycological Research 105(7): 773–782. <https://doi.org/10.1017/S0953756201004257>
- Wang XW, Liu SL, Zhou LW (2023) An updated taxonomic framework of Hymenochaetales (Agaricomycetes, Basidiomycota). Mycosphere 14(1): 452–496. <https://doi.org/10.5943/mycosphere/14/1/6>
- White TJ, Bruns TD, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols, a guide to methods and applications. Academic, San Diego, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wu F, Zhou LW, Yang ZL, Bau T, Li TH, Dai YC (2019) Resource diversity of Chinese macrofungi: Edible, medicinal and poisonous species. Fungal Diversity 98(1): 1–76. <https://doi.org/10.1007/s13225-019-00432-7>
- Wu F, Man XW, Tohtirjap A, Dai YC (2022a) A comparison of polypore funga and species composition in forest ecosystems of China, North America, and Europe. Forest Ecosystems 9: 100051. <https://doi.org/10.1016/j.fecs.2022.100051>
- Wu F, Zhou LW, Vlasák J, Dai YC (2022b) Global diversity and systematics of Hymenochaetaceae with poroid hymenophore. Fungal Diversity 113(1): 1–192. <https://doi.org/10.1007/s13225-021-00496-4>
- Yang Y, Xu Y, Wang L, Jiang QQ, Su JQ, Li R, Zhou HM, Zhao CL (2025) Multigene phylogeny of seven wood-inhabiting fungal orders in Basidiomycota, and proposal of a new genus and thirteen new species. Mycosphere 16(1): 245–295. <https://doi.org/10.5943/mycosphere/16/1/4>

- Yuan Y, Bian LS, Wu YD, Chen JJ, Wu F, Liu HG, Zeng GY, Dai YC (2023) Species diversity of pathogenic wood-rotting fungi (Agaricomycetes, Basidiomycota) in China. *Mycology* 14: 204–226. <https://doi.org/10.1080/21501203.2023.2238779>
- Zhang X, Shi LY, Xie XJ, Wei TZ, Guo YB, Zhao SY, Li SM, Li GJ (2023) Biological characteristics and domestication of *Phellinus orientoasiaticus*. *Mycosystema* 42(4): 883–895. <https://doi.org/10.13346/j.mycosystema.220522>
- Zhang SY, Yu SW, Wei YL (2024) A comparative study on the bioactive components and antioxidant activities of *Phellinus orientoasiaticus* and two species of *Sanghuangporus*. *Mycosystema* 43(2): 230220. <https://doi.org/10.13346/j.mycosystema.230220>
- Zhao H, Wu F, Maurice S, Pavlov IN, Krutovsky KV, Liu HG, Yuan Y, Dai YC (2024a) Large-scale phylogenomic insights into the evolution of the Hymenochaetales. *Mycology*: 1–18. <https://doi.org/10.1080/21501203.2024.2391527>
- Zhao H, Wu YD, Yang ZR, Liu HG, Wu F, Dai YC, Yuan Y (2024b) Polypore funga and species diversity in tropical forest ecosystems of Africa, America and Asia, and a comparison with temperate and boreal regions of the Northern Hemisphere. *Forest Ecosystems* 11: 100200. <https://doi.org/10.1016/j.fecs.2024.100200>
- Zhou LW, Xue HJ (2012) *Fomitiporia pentaphylacis* and *F. tenuitubus* spp. nov. (Hymenochaetales, Basidiomycota) from Guangxi, southern China. *Mycological Progress* 11(4): 907–913. <https://doi.org/10.1007/s11557-012-0806-1>
- Zhou M, Dai YC, Vlasák J, Liu HG, Yuan Y (2023) Updated systematics of *Trichaptum* s.l. (Hymenochaetales, Basidiomycota). *Mycosphere* 14: 815–917. <https://doi.org/10.5943/mycosphere/14/1/11>